

Oxidant Gases

by Michael J. Evans*†

The acute and chronic action of the oxidant gases ozone, nitrogen dioxide and oxygen on the morphological appearance of cells of the alveolar and bronchiolar epithelium is reviewed. Type I cells of the alveolar and ciliated cells of the bronchiolar epithelium appear to be sensitive targets for the oxidant gases. The degree of damage is influenced by age, nutritional status and the development of tolerance.

Introduction

The airways of the lung are lined with a layer of ciliated and nonciliated cells (1). In the large airways (trachea, main bronchi, and bronchi), the nonciliated cell population comprises serous, goblet, intermediate, and basal cells. In the small airways (terminal and respiratory bronchioles), Clara cells make up the nonciliated cell population. Goblet, serous, and Clara cells usually protrude above adjacent ciliated cells and contain secretory granules. In the alveoli, large, flat squamous cells (Type I) cover about 97% of the alveolar surface (2). The remaining 3% of the gas-exchanging area is covered by cuboidal cells (Type II), which usually contain secretory granules (lamellar bodies). Beneath the epithelial surface is the interstitium, which is composed of acellular supporting materials (collagen and elastin), fibroblasts, and migratory cells. Within the interstitium there is a vast network of capillaries lined with endothelial cells. Together, the Type I cell, interstitium, and endothelial cell make up the air-blood barrier of the lung. Macrophages are present in the surface-active film covering the alveoli. This alveolar lining layer is continuous with a mucoid layer lining the airways.

The purpose of this paper is to describe the effects of oxidant gases on these structures. When animals are exposed to oxidant gases such as nitrogen dioxide (NO_2), ozone (O_3) or oxygen (O_2), the gas first contacts the acellular lining layer, then the macrophages, epithelial cells, and the interstitium, and finally the endothelium and capillary blood of the alveoli. The ability of these gases to affect cells of the lung is influenced by their concentration, aerodynamics, solubility, reactivity, and dilution as the air permeates the more distal alveoli (3-5). The ability of the animal to resist oxidant damage depends on its age and nutritional status and the degree

to which it has adapted or become tolerant to the gas (6-9). Oxidant-induced injury to lung cells is discussed in this paper with respect to these factors.

Acute Lung Injury

When animals are exposed continuously to sublethal concentrations of NO_2 (2 to 17 ppm) or to O_3 (0.5 to 1.0 ppm), the epithelial cells lining the airways and alveoli are damaged (5,10-20). The primary anatomical site of cellular damage is the alveoli in the central region of the pulmonary acinus and the adjacent bronchioles (terminal or respiratory) (5,18). The peripheral portion of the pulmonary acinus and larger airways generally are not affected as severely (Figs. 1 and 2). Morphologically, greater injury is expressed as a large amount of alveolar damage and more involvement of the upper airways (3,21). These characteristics are thought to be due to a concentration gradient that exists during inhalation of NO_2 or O_3 .

In the bronchiolar regions, the first morphological changes are observed in ciliated cells (5,14,17,19,22-24). By 4 hr of exposure, a loss of cilia begins. By 8 hr, the




FIGURE 1. Tissue architecture in a control rat. $\times 55$.

*SRI International, 333 Ravenswood Avenue, Menlo Park, CA 94025.

†Present address: Inhalation Toxicology Research Institute, Lovelace Biomedical and Environmental Research Institute, P. O. Box 5890, Albuquerque, NM 87185.




FIGURE 2. Area similar to that in Figure 1 after 48 hr exposure to 17 ppm NO_2 . The increased density of alveoli near the opening of terminal bronchioles is caused by proliferation of cells and accumulation of macrophages associated with the damage caused during the first 24 hr of exposure and its repair. $\times 55$.




FIGURE 3. Micrograph of portion of bronchiolar epithelium from a control rat, demonstrating a Clara cell protruding above adjacent cells. $\times 1250$.

loss is obvious, and sloughing of dying and necrotic ciliated cells is seen (Figs. 3–5). After 16 hr of exposure, the ciliated and nonciliated cells of the bronchiolar epithelium become uniform in height. Although Clara cells are relatively resistant to injury, they lose secretory granules during the first 24 hr of exposure (25,26). By the second day of exposure, cellular hypertrophy and hyperplasia are evident in the bronchiolar epithelium and foci of multilayered cells become obvious (13,26–29). From day 2 through 15, the epithelium tends to maintain this cellular appearance, with a distinct absence of cilia (Fig. 6). During this period, numerous ciliated vacuoles are observed (Fig. 7), along with




FIGURE 4. Bronchiolar epithelium after 24 hr exposure to 15 ppm NO_2 . Note the lack of cilia. $\times 1250$.




FIGURE 5. Electron micrograph of bronchiolar epithelium after 8 hr of exposure to 14 ppm NO_2 , demonstrating necrotic cells that had sloughed off the bronchiolar wall. $\times 5000$.

rodlike structures in nonciliated cells (Fig. 8). When the animals are allowed to recover in air, normal ciliated, and nonciliated cells appear again by day 7 of recovery (Fig. 9) (10,25,30–33).

In alveolar regions, the first morphological changes are observed also by the fourth hour of exposure




FIGURE 6. Bronchiolar epithelium after 15 days of exposure to 17 ppm NO_2 . $\times 1250$.




FIGURE 7. Ciliated vacuole in a ciliated cell after 15 days of exposure to 12 ppm NO_2 . $\times 3710$.




FIGURE 8. Rodlike structures in nonciliated cells after 5 days of exposure to 12 ppm NO_2 . $\times 5822$.



FIGURE 9. Bronchiolar epithelium in a rat exposed to 15 ppm NO_2 for 24 hr and allowed to recover in air for 7 days. Compare with Fig. 3 and note the return of a normal ciliary bed. $\times 1250$.

(18,19,34,35). At this time, Type I cells near the opening of the terminal bronchioles exhibit swelling, and occasional areas of alveoli have lost the covering Type I cells, leaving basement membrane exposed (Figs. 10 and 11). These changes increase in intensity so that, after a day of continuous exposure, the larger areas of alveoli have lost their damaged Type I cells (Fig. 12). Cellular debris and some interstitial and alveolar edema become apparent (10,15-19,22,35). Type II cells appear to be relatively resistant to NO_2 or O_3 , as do alveolar macrophages. Endothelial cells appear not to be affected by NO_2 , but occasionally swelling has been reported during exposure to O_3 .

After 48 hr of continuous exposure, areas previously devoid of Type I cells are covered by a low cuboidal epithelium (Figs. 2, 13, and 14). These cells are derived




FIGURE 10. Example of injured alveoli near the opening of the terminal bronchiole after 12 hr of exposure to 15 ppm NO_2 . $\times 555$.

from Type II cells but lack lamellar bodies (36-39). Cellular debris, mucoid material and macrophages are present near the openings of the terminal and respiratory bronchioles. By day 4 of exposure, the epithelium has assumed a more squamous appearance, and most of the cellular and amorphous debris has been eliminated (Fig. 15). However, the number of epithelial cells present has increased and the air-blood barrier is thicker (3,10,16,40,41).

Other cells in the alveoli also multiply during exposure to NO_2 or O_3 (38,39,42). The number of dividing macrophages increases by day 3 of exposure. Whether this is in response to the presence of cellular debris or to hormonal stimulation associated with Type II cell division is not clear. The number of interstitial cells and the thickness of the interstitial space also increased (20,33,43-47). In the capillaries, there is often an increase in the number of circulating mononuclear cells. The tissue remains in this condition for at least an additional 15 days (Fig. 16). If animals are allowed to recover in air, the epithelium soon assumes a normal appearance (10,26). These changes have been described in a variety of animals, including mice, rats, rabbits, cats, dogs and primates.




FIGURE 11. Swelling and sloughing of Type I cells after 12 hr exposure to 15 ppm NO_2 . $\times 7000$.




FIGURE 12. Necrotic Type I cell sloughing off the alveolar wall after 24 hr of exposure to 14 ppm NO_2 . Arrow indicates area of basement membrane. $\times 6700$.




FIGURE 13. Cuboidal epithelium and cell debris in alveoli near the opening of a terminal bronchiole. $\times 275$.

From these studies, three important conclusions were derived. First, the cells most vulnerable to NO_2 or O_3 (target cells) are the ciliated bronchiolar and Type I alveolar cells. Both cell types have a much larger surface area exposed to the NO_2 than do the more resistant types (Type II cells and Clara cells). Second, the epithelium can repair itself in the presence of sublethal levels of NO_2 or O_3 . Third, the new cells are relatively resistant to continuous exposure to the same concentration of NO_2 or O_3 . Because repair does occur and the new cells are resistant to injury, it is important to understand the time sequence of injury when studying the effects of NO_2 or O_3 on these target cells.

When animals are exposed continuously to lethal (100%) and sublethal (85%) concentrations of O_3 , damage is observed in the airways and alveoli. However, there is no preferred anatomical site of damage, and damage is observed diffusely throughout the lung (48-57).

In the airways (bronchi and bronchioles), swelling of nonciliated cells and loss of cilia are observed after 2 days of exposure (58,59). By day 3, necrotic nonciliated cells are present and sloughing of ciliated cells is




FIGURE 14. Electron micrograph of the cuboidal epithelium that results from proliferation of Type II cells for renewal of injured Type I epithelium. $\times 4000$.



FIGURE 15. Tissue architecture after 4 days of exposure to 17 ppm NO₂, demonstrating the return of the epithelium to a more squamous type. $\times 55$.

observed. Proliferation of cells to repair such injury begins by day 4 if the animals survive the exposure (29,60).

In the alveoli, slight swelling is observed in endothelial cells on day 2 of exposure. By day 3 of exposure, individual endothelial cells undergo cytoplasmic changes and fragmentation, which can result in destruction of up to 50% of the capillary bed (Fig. 17) (54-56,61). These changes are accompanied by increased thickness of the interstitium and damage to Type I cells. Between day 3 and day 4, a variety of lesions—such as fibrinous exudation, hyaline membrane formation and capillary congestion—are observed. In animals surviving these effects, proliferation of Type II cells, similar to that described for NO₂ and O₃ exposure, begins between day 4 and day 5 (36,38,49,62). Surviving animals allowed to recover in air for 56 days have a nearly normal epithelial lining, but septal scars remain due to the increase in interstitial tissue (54).

If animals are exposed to sublethal concentrations of O₂ (85%), changes in the endothelium similar to those seen with 100% O₂ occur (3,61,63,64). However, very little

damage is observed in the Type I epithelium. There is a net increase in Type II cells and a large increase in interstitial cells between day 5 and day 14 of exposure.




FIGURE 16. Tissue architecture after 15 days of exposure to 17 ppm NO_2 . $\times 46$.

Some investigators consider these changes to be adaptive (61,65). Results similar to these have been described in many species of animals, including man (53).

These studies suggest that the morphological target cells during exposure to O_2 are ciliated and nonciliated cells in the airways and endothelial cells in the alveoli. Although other cells in the alveoli may be affected during exposure, it is not clear whether this is due to the oxygen or is a secondary response to the initial injury.

Factors That Affect the Amount of Tissue Injury

The target cells associated with damage to the airways and alveoli during exposure to NO_2 , O_3 or O_2 were described in the previous section. The degree to which target cells in the tissue are affected is subject to several factors. Obviously, concentration affects the amount of tissue damage. However, age, nutritional status, and the development of tolerance are also important in this regard.




FIGURE 17. Damage to endothelial cell in a rat after 2 days of exposure to 100% O_2 . $\times 11,600$.

A study of aging (10,30) revealed that relatively old rats exposed to 14 ppm NO_2 suffered greater injury than did young rats exposed to 17 ppm NO_2 . Morphologically, the nature of the injury observed in the airways and alveoli was the same for both groups, but the area of damage in the alveoli was more extensive in the older animals and was associated with increased alveolar edema (10,30). Further studies showed that initiation of repair of damaged epithelium in older animals was delayed by approximately 1 day. Both factors—the greater extent of tissue damage and the retarded onset of cell proliferation—are likely to be responsible for the edema and hence the higher mortality rate in the older animals. After exposure to about 14 ppm NO_2 , 12% of 11-month-old rats died, but 35% of the 19-month-old rats died. It is interesting, however, that older animals that survived the first 3 days of exposure were able to repair the damage, as young rats did, despite continuous exposure to NO_2 . The reason for the difference in initial susceptibility to injury was not determined. Older animals exposed to O_3 or O_2 are also known to be more susceptible to injury (61,66).

In contrast, Stephens et al. (67) observed resistance to high concentrations of NO_2 or O_3 in rats exposed from birth until they were about 30 days old. In this study, loss of cilia was reported in the terminal bronchioles of nursing rats from 8 to 20 days of age. In the alveoli, no evidence of damage to Type I cells was seen until the rats were 21 to 35 days old. The basis for the refractory nature of the alveolar epithelium up to 20 days was not apparent. Young animals also are more resistant to the toxic effects of O_2 (68–70).

Another factor that may affect oxidant damage is the nutritional status of the animal. For example, the tissue levels of the natural antioxidants vitamin E and selenium are important mediators of cell damage. In a series of studies with rats fed diets containing various amounts of vitamin E prior to exposure to NO_2 , Fletcher and Tappel (71) found that vitamin E-supplemented animals died later than those fed vitamin E-deficient diets. Death was not averted, however, in animals supplemented with vitamin E during exposure to NO_2 .

Recently, Evans et al. (72) demonstrated that rats depleted of vitamin E and selenium suffered more Type I cell damage in response to NO_2 exposure than did those fed adequate levels of these compounds. However, the supplemented groups were protected for only 6 to 12 hr. After this time, all groups incurred the same amount of damage. Chow et al. (73) reported more injury in rats depleted of vitamin E than in rats on supplemented diets when they were exposed to O_3 . These studies demonstrate that biological antioxidants play a role in affecting the extent of damage sustained by alveolar epithelium during exposure to NO_2 or O_3 .

Although NO_2 and O_3 injure pulmonary cells of rats upon initial exposure, the tissue becomes relatively tolerant to the same concentration during continuous exposure. If the animals are removed from the NO_2 - or

O_3 -containing environment, the tolerance decreases. Using intermittent exposures, Creasia et al. (52) showed that tolerance that had developed to 10 ppm NO_2 persisted for about 7 days before re-exposure to the same concentration again caused significant injury. Similar results were obtained from rats exposed to ozone (38,74). Animals exposed to 85% O_2 also developed tolerance after 5 days of exposures (3,61,64,75–78).

Effects of Chronic Exposure

Early pathologic events during and after limited initial exposure of rats to NO_2 or O_3 are reparable, as described, but persistent exposure to subacute concentrations of at least 14 ppm NO_2 for several months leads to permanent distortion of the architecture of the rats' lungs (Figs. 18 and 19), with loss of alveolar tissue (27,28,31,43,79,80,81,82). Lower concentrations also affect the lungs, but 1 or 2 ppm—even for the lifetime of the rat—fails to induce parenchymal destruction (82), although respiratory rates and lung volumes become elevated and metabolic changes are seen in the bronchial epithelium (43).




FIGURE 18. Parenchyma of a 24-month-old control rat. $\times 20$.




FIGURE 19. Parenchyma of a rat exposed for prolonged, varying periods to ~ 15 ppm NO_2 over a 24-month period, exhibiting a loss of alveolar surface area. $\times 20$.

In chronically exposed animals, the characteristic paucity of cilia in the airways and the relatively uniform hypertrophy of the epithelium of the small airways, with their occasional foci of hyperplasia, indicate a much reduced—but continuing—response of the lining cells to the persistent presence of NO_2 or O_3 . From the onset of exposure, the most severely affected tissues of the airways and parenchyma—mainly the terminal bronchioles, alveolar ducts and adjacent alveoli—are infiltrated with inflammatory cells. These are largely macrophages both within the alveoli and the small airways (where they often fill the former close to the small airways and ducts) and in the interstitial tissue. Other types of mononuclear cells, and occasionally granulocytic cells, are included. Mast cells are often seen dispersed throughout the lung.

The ubiquitous network of connective tissue elements that permeates the alveolar walls and small airways reveals changes in staining properties of both the elastic tissue and the collagen strands in long-exposed rats (43). Besides appearing to become thicker with time, the elastic strands tend to fracture and some of the collagen fibers appear as thick, ropelike structures (47). Simultaneously, the basement membranes supporting both the alveolar epithelium and the capillary endothelium become broader in proximal alveoli, in contrast with the unusually thin, attenuated alveolar walls of more peripheral alveoli. In general, pulmonary blood volume per gram of tissue is reduced in the heavy, voluminous, nonedematous lungs that develop during exposure to high, subacute concentrations of about 15 ppm NO_2 (28).

The internal diameters of terminal bronchioles decrease during prolonged exposure as their walls (composed mainly of smooth muscle, connective tissue, and epithelial lining) become thicker (33). The diameters are reduced further because of an accumulation of mucous and serous secretions from the increased number of peripherally activated "goblet" and other secreting cells of the bronchial and bronchiolar mucosa; hence, airflow is correspondingly reduced. Cellular debris, fibrinous exudate, and the attendant macrophages and other cells of the immune system are additional obstacles to flow.

Although these changes do not involve target cell toxicity, they do reflect changes in the tissue due to chronic exposure. Whether these changes reflect reparative or adaptive phenomena or the direct action of NO_2 or O_3 on the structural organization of the lung is not clear. However, because of the damage, long-term studies should be included in any research on oxidant-induced lung toxicity.

I am grateful to colleagues listed among the references for their basic contributions and to many other collaborators for excellent assistance.

This work was supported by National Institutes of Health Grant No. ES00842.

REFERENCES

1. Breeze, R. G., and Wheeldon, E. G. The cells of the pulmonary airways. *Am. Rev. Respir. Dis.* 116: 705-777 (1977).
2. Weibel, E. R., Gehr, P., Haies, D., Gil, J., and Bachofen, M. The cell population of the normal lung. In: *Lung Cells in Disease* (A. Bouhuys, Ed.), Elsevier/North-Holland Biomedical Press, Amsterdam, Holland, 1976, pp. 3-16.
3. Crapo, J. D., March-Salin, J., Ingram, P., and Pratt, P. C. Tolerance and cross-tolerance using NO_2 and O_3 . II. Pulmonary morphology and morphometry. *J. Appl. Physiol.* 44: 370-379 (1978).
4. Evans, M. J., and Freeman, G. Morphological and pathological effects of NO_2 on the rat lung. In: *Nitrogen Oxides and Their Effects on Health* (S. D. Lee, Ed.), Ann Arbor Science Pubs., Inc., Ann Arbor, Michigan, 1980, pp. 243-265.
5. Plopper, C. G., Dungworth, D. L., and Tyler, W. S. Pulmonary lesions in rats exposed to ozone. *Am. J. Pathol.* 71: 375-386 (1973).
6. Clark, J. M., and Lambertson, C. J. Pulmonary oxygen toxicity: a review. *Pharmacol. Rev.* 23: 37-133 (1971).
7. Haugaard, N. Cellular mechanisms of oxygen toxicity. *Physiol. Rev.* 48: 311-373 (1968).
8. Mustafa, M. G., and Tierney, D. F. Biochemical and metabolic changes in the lung with oxygen, ozone, and nitrogen dioxide toxicity. *Am. Rev. Respir. Dis.* 118: 1061-1090 (1978).
9. Stokinger, H. E. Ozone toxicology. *Arch. Environ. Health* 10: 719-731 (1965).
10. Cabral-Anderson, L. J., Evans, M. J., and Freeman, G. Effects of NO_2 on the lungs of aging rats. I. Morphology. *Exptl. Mol. Pathol.* 27: 353-365 (1977).
11. Castleman, W. L., Tyler, W. S., and Dungworth, D. L. Lesions in respiratory bronchioles and conducting airways of monkeys exposed to ambient levels of ozone. *Exptl. Mol. Pathol.* 26: 384-400 (1977).
12. Evans, M. J. Cell death and cell renewal in small airways and alveoli. In: *Mechanisms in Respiratory Toxicology* (H. Witschi and P. Nettesheim, Eds.), CRC Press, Cleveland, Ohio, 1981.
13. Kleinerman, J. Effects of NO_2 in hamsters: autoradiographic and electron microscopic aspects. In: *Inhalation Carcinogenesis* (M. G. Hanna, Jr., P. Nettesheim and J. R. Gilbert, Eds.), U.S. Atomic Energy Commission Division of Technical Information, Oak Ridge, TN, 1970, p. 271.
14. Mellick, P. W., Dungworth, D. L., Schwartz, L. W., and Tyler, W. S. Short-term morphologic effects of high ambient levels of ozone on lungs of rhesus monkeys. *Lab. Invest.* 36: 82-90 (1977).
15. Plopper, C. G., Dungworth, D. L., and Tyler, W. S. Morphometric evaluation of pulmonary lesions in rats exposed to ozone. *Am. J. Pathol.* 71: 395-408 (1973).
16. Plopper, C. G., Dungworth, D. L., Tyler, W. S., and Chow, C. K. Pulmonary alterations in rats exposed to 0.2 and 0.1 ppm ozone: a correlated morphological and biochemical study. *Arch. Environ. Health* 34: 390-395 (1979).
17. Schwartz, L. W., Dungworth, D. L., Mustafa, M. G., Tarkington, B. K., and Tyler, W. S. Pulmonary responses of rats to ambient levels of ozone. Effects of 7-day intermittent or continuous exposure. *Lab. Invest.* 34: 565-578 (1976).
18. Stephens, R. J., Freeman, G., and Evans, M. J. Early response of lungs to low levels of nitrogen dioxide. *Arch. Environ. Health* 24: 160-179 (1972).
19. Stephens, R. J., Sloan, M. F., Evans, M. J., and Freeman, G. Early response of lung to low levels of ozone. *Am. J. Pathol.* 74: 31-58 (1974).
20. Werthamer, S., Schwarz, L. H., Carr, J. J., and Soskind, L. Ozone-induced pulmonary lesions. *Arch. Environ. Health* 20: 16-21 (1970).
21. Evans, M. J., Dekker, N. P., Cabral-Anderson, L. J., and Freeman, G. Quantitation of damage to the alveolar epithelium by means of Type 2 cell proliferation. *Am. Rev. Respir. Dis.* 118: 787-790 (1978).
22. Boatman, E. S., Sato, S., and Frank, R. Acute effects of ozone on

- cat lungs. II. Structural. *Am. Rev. Respir. Dis.* 110: 157-169 (1974).
23. Castleman, W. L., Dungworth, D. L., Schwartz, L. W., and Tyler, W. S. Acute respiratory bronchiolitis. *Am. J. Pathol.* 98: 811-840 (1980).
24. Ibrahim, A. L., Zee, Y. C., and Osebold, J. W. The effects of ozone on the respiratory epithelium of mice. II. Ultrastructural alterations. *J. Environ. Pathol. Toxicol.* 3: 251-258 (1980).
25. Evans, M. J., Cabral-Anderson, L. J., and Freeman, G. Role of the Clara cell in renewal of the bronchiolar epithelium. *Lab. Invest.* 38: 648-655 (1978).
26. Evans, M. J., Johnson, L. V., Stephens, R. J., and Freeman, G. Renewal of terminal bronchiolar epithelium following exposure to O_3 and NO_2 . *Lab. Invest.* 35: 246-257 (1976).
27. Freeman, G., Crane, S. C., Stephens, R. J., and Furiosi, N. J. Pathogenesis of the nitrogen dioxide-induced lesion in the rat lung: A review and presentation of new observations. *Am. Rev. Respir. Dis.* 98: 429-443 (1968).
28. Freeman, G., Crane, S. C., Stephens, R. J., and Furiosi, N. J. The subacute nitrogen dioxide-induced lesion in the rat lung. *Arch. Environ. Health* 18: 609-612 (1969).
29. Lum, H., Schwartz, L. W., Dungworth, D. L., and Tyler, W. S. A comparative study of cell renewal after exposure to ozone or oxygen. Response of terminal bronchiolar epithelium in the rat. *Am. Rev. Respir. Dis.* 118: 335-345 (1978).
30. Evans, M. J., Cabral-Anderson, L. J., and Freeman, G. Effects of NO_2 on the lungs of aging rats. II. Cell proliferation. *Exptl. Mol. Pathol.* 27: 366-376 (1977).
31. Freeman, G., Crane, S. C., Furiosi, N. J., Stephens, R. J., Evans, M. J., and Moore, W. D. Covert reduction in ventilatory surface in rats during prolonged exposure to subacute nitrogen dioxide. *Am. Rev. Respir. Dis.* 106: 563-579 (1972).
32. Stephens, R. J., Freeman, G., Crane, S. C., and Furiosi, N. J. Ultrastructural changes in the terminal bronchioles of the rat during continuous, low-level exposure to NO_2 (ciliogenesis). *Exptl. Mol. Pathol.* 14: 1-19 (1971).
33. Stephens, R. J., Freeman, G., Stara, J. F., and Coffin, D. L. Cytologic changes in dog lungs induced by chronic exposure to ozone. *Am. J. Pathol.* 73: 711-726 (1973).
34. Stephens, R. J., Evans, M. J., Sloan, M. F., and Freeman, G. A comprehensive ultrastructural study of pulmonary injury and repair in the rat resulting from exposures to less than one ppm ozone. *Chest* 65: 11S-13S (1974).
35. Stephens, R. J., Sloan, M. F., Evans, M. J., and Freeman, G. Alveolar Type 1 cell response to exposure to 0.5 ppm O_3 for short periods. *Exptl. Mol. Pathol.* 20: 11-23 (1974).
36. Evans, M. J., Cabral, L. J., Stephens, R. J., and Freeman, G. Renewal of alveolar epithelium in the rat following exposure to NO_2 . *Am. J. Pathol.* 70: 171-194 (1973).
37. Evans, M. J., Cabral, L. J., Stephens, R. J., and Freeman, G. Transformation of alveolar Type 2 cells to Type 1 cells following exposure to NO_2 . *Exptl. Mol. Pathol.* 22: 142-150 (1973).
38. Evans, M. J., Johnson, L. V., Stephens, R. J., and Freeman, G. Cell renewal in the lungs of rats exposed to low levels of ozone. *Exptl. Mol. Pathol.* 24: 70-83 (1976).
39. Evans, M. J., Stephens, R. J., Cabral, L. J., and Freeman, G. Cell renewal in the lungs of rats exposed to low levels of NO_2 . *Arch. Environ. Health* 24: 180-188 (1972).
40. Boorman, G. A., Schwartz, L. W., and Dungworth, D. L. Pulmonary effects of prolonged ozone insult in rats. Morphometric evaluation of the central acinus. *Lab. Invest.* 43: 108-115 (1980).
41. Yuen, T. G. A., and Sherwin, R. P. Hyperplasia of Type II pneumocytes and nitrogen dioxide (10 ppm): a quantitation based on electron photomicrographs. *Arch. Environ. Health* 24: 178-188 (1971).
42. Evans, M. J., Cabral, L. J., Stephens, R. J., and Freeman, G. Cell division of alveolar macrophages in the rat following exposure to NO_2 . *Am. J. Pathol.* 70: 195-203 (1973).
43. Freeman, G., Juhos, L. T., Furiosi, N. J., Mussenden, R., Stephens, R. J., and Evans, M. J. Pathology of pulmonary disease from exposure to interdependent ambient gases (nitrogen dioxide and ozone). *Arch. Environ. Health* 29: 203-210 (1974).
44. Last, J. A., and Greenberg, D. B. Ozone-induced alterations in collagen metabolism of rat lungs. II. Long-term exposures. *Toxicol. Appl. Pharmacol.* 55: 108-114 (1980).
45. Last, J. A., Greenberg, D. B., and Castleman, W. L. Ozone-induced alterations in collagen metabolism of rat lungs. *Toxicol. Appl. Pharmacol.* 51: 247-258 (1979).
46. Penha, P. D., and Werthamer, S. Pulmonary lesions induced by long-term exposure to ozone. *Arch. Environ. Health* 29: 282-289 (1974).
47. Stephens, R. J., Freeman, G., and Evans, M. J. Ultrastructural changes in connective tissue in lungs of rats exposed to NO_2 . *Arch. Intern. Med.* 127: 873-883 (1971).
48. Adamson, I. Y. R., Bowden, D. H., and Wyatt, J. P. Oxygen poisoning in mice. Ultrastructural and surfactant studies during exposure and recovery. *Arch. Pathol.* 90: 463-472 (1970).
49. Bowden, D. H., and Adamson, I. Y. R. Reparative changes following pulmonary cell injury. *Arch. Pathol.* 92: 279-283 (1971).
50. Bowden, D. H., Adamson, I. Y. R., and Wyatt, J. P. Reaction of the lung cells to a high concentration of oxygen. *Arch. Pathol.* 86: 671-675 (1968).
51. Coalson, J. J., Beller, J. J., and Greenfield, L. J. Effects of 100 per cent oxygen ventilation on pulmonary ultrastructure and mechanics. *J. Pathol.* 104: 267-273 (1971).
52. Creasia, D. A., Nettesheim, P., and Kim, J. C. S. Stimulation of DNA synthesis in the lungs of hamsters exposed intermittently to nitrogen dioxide. *J. Toxicol. Environ. Health* 2: 1173-1181 (1977).
53. Gould, V. E., Tosco, R., Wheelis, R. F., Gould, N. S., and Kapanci, Y. Oxygen pneumonitis in man. *Lab. Invest.* 26: 499-508 (1972).
54. Kapanci, Y., Weibel, E. R., Kaplan, H. P., and Robinson, F. R. Pathogenesis and reversibility of the pulmonary lesions of oxygen toxicity in monkeys. II. Ultrastructural and morphometric studies. *Lab. Invest.* 20: 101-118 (1969).
55. Kaplan, H. P., Robinson, F. R., Kapanci, Y., and Weibel, E. R. Pathogenesis and reversibility of the pulmonary lesions of oxygen toxicity in monkeys. I. Clinical and light microscope studies. *Lab. Invest.* 20: 94-100 (1969).
56. Kistler, G. S., Caldwell, P. R. B., and Weibel, E. R. Development of fine structural damage to alveolar and capillary lining cells in oxygen-poisoned rat lungs. *J. Cell Biol.* 33: 605-628 (1967).
57. Schaffner, F., Felig, P., and Trachtenberg, E. Structure of rat lung after protracted oxygen breathing. *Arch. Pathol.* 83: 99-107 (1967).
58. Ludwin, S. K., Northway, W. H., Jr., and Bensch, K. G. Oxygen toxicity in the newborn. Necrotizing bronchiolitis in mice exposed to 100% oxygen. *Lab. Invest.* 31: 425-435 (1974).
59. Obora, H., Sekimoto, M., and Iwai, S. Alterations to the bronchial and bronchiolar surfaces of adult mice after exposure to high concentrations of oxygen. *Thorax* 34: 479-485 (1979).
60. Evans, M. J., Hackney, J. D., and Bils, R. F. Effects of a high concentration of oxygen on cell renewal in the pulmonary alveoli. *Aerosp. Med.* 40: 1365-1368 (1969).
61. Crapo, J. D., Barry, B. E., Foscue, H. A., and Shelburne, J. Structural and biochemical changes in rat lungs occurring during exposures to lethal and adaptive doses of oxygen. *Am. Rev. Respir. Dis.* 122: 123-142 (1980).
62. Adamson, I. Y. R., and Bowden, D. H. Derivation of Type 1 epithelium from Type 2 cells in the developing rat lung. *Lab. Invest.* 32: 736-745 (1975).
63. Crapo, J. D., Peters-Golden, M., Marsh-Salin, J., and Shelburne, J. S. Pathologic changes in the lungs of oxygen-adapted rats. A morphometric analysis. *Lab. Invest.* 39: 640-653 (1978).
64. Crapo, J. D., Sjoström, K., and Drew, R. T. Tolerance and cross-tolerance using NO_2 and O_3 . I. Toxicology and biochemistry. *J. Appl. Physiol.* 44: 364-369 (1970).
65. Yamamoto, E., Wittner, M., and Rosenbaum, R. M. Resistance and susceptibility to oxygen toxicity by cell types of the gas-blood barrier of the rat lung. *Am. J. Pathol.* 59: 409-435 (1970).
66. Evans, M. J., Mayr, W., Bils, R. F., and Loosli, C. G. Effects of

- ozone on cell renewal in pulmonary alveoli of aging mice. *Arch. Environ. Health* 22: 450-453 (1971).
67. Stephens, R. J., Sloan, M. F., Groth, D. G., Negi, D. S., and Lunan, K. D. Cytologic response of postnatal rat lungs to O₃ or NO₂ exposure. *Am. J. Pathol.* 93: 183-200 (1978).
68. Anderson, W. R., Strickland, M. B., Tsai, S. H., and Haglin, J. J. Light microscopic and ultrastructural study of the adverse effects of oxygen therapy on the neonate lung. *Am. J. Pathol.* 73: 327-339 (1973).
69. Bonikos, D. S., Bensch, K. G., Ludwin, S. K., and Northway, W. H. Oxygen toxicity in the newborn. The effect of prolonged 100 per cent O₂ exposure on the lungs of newborn mice. *Lab. Invest.* 32: 619-635 (1975).
70. Frank, L., Bucher, J. R., and Roberts, R. J. Oxygen toxicity in neonatal and adult animals of various species. *J. Appl. Physiol.* 45: 699-704 (1978).
71. Fletcher, B. L., and Tappel, A. L. Protective effects of dietary α -tocopherol in rats exposed to toxic levels of ozone and nitrogen dioxide. *Environ. Res.* 6: 165-175 (1973).
72. Evans, M. J., Cabral-Anderson, L. J., Dekker, N. P., and Freeman, G. The effects of dietary antioxidants on NO₂-induced injury to Type 1 alveolar cells. *Chest* 80: 5-8 (1980).
73. Chow, C. K., Plopper, C. G., and Dungworth, D. L. Influence of dietary vitamin E on the lungs of ozone-exposed rats. *Environ. Res.* 20: 309-317 (1979).
74. Chow, C. K., Hussain, M. Z., Cross, C. E., Dungworth, D. L., and Mustafa, M. G. Effects of low levels of ozone on rat lungs. I. Biochemical responses during recovery and reexposure. *Exptl. Mol. Pathol.* 25: 182-188 (1976).
75. Crapo, J. D., and Tierney, D. F. Superoxide dismutase and pulmonary oxygen toxicity. *Am. J. Physiol.* 226: 1401-1407 (1974).
76. Frank, L., Yam, J., and Roberts, R. J. The role of endotoxin in protection of adult rats from oxygen-induced lung toxicity. *J. Clin. Invest.* 61: 269-275 (1978).
77. Yam, J., Frank, L., and Roberts, R. J. Oxygen toxicity: Comparison of lung biochemical responses in neonatal and adult rats. *Pediatr. Res.* 12: 115-119 (1978).
78. Yam, J., and Roberts, R. J. Pharmacological alteration of oxygen-induced lung toxicity. *Toxicol. Appl. Pharmacol.* 47: 367-376 (1979).
79. Freeman, G., Furiosi, N. J., and Haydon, G. B. Effects of continuous exposure of 0.8 ppm NO₂ on respiration of rats. *Arch. Environ. Health* 13: 454-456 (1966).
80. Freeman, G., and Haydon, G. B. Emphysema after low-level exposure to NO₂. *Arch. Environ. Health* 8: 125-128 (1964).
81. Freeman, G., Stephens, R. J., Coffin, D. L., and Stara, J. F. Changes in dogs' lungs after long-term exposure to ozone. *Arch. Environ. Health* 26: 209-216 (1973).
82. Freeman, G., Stephens, R. J., and Furiosi, N. J. Lesion of the lung in rats continuously exposed to two parts per million of nitrogen dioxide. *Arch. Environ. Health* 17: 181-192 (1968).
83. Juhos, L. T., Green, D. P., Furiosi, N. J., and Freeman, G. A quantitative study of stenosis in the respiratory bronchiole of the rat in NO₂-induced emphysema. *Am. Rev. Respir. Dis.* 121: 541-549 (1980).